

The Compton effect in strongly magnetized plasma

M. V. Chistyakov, D. A. Romyantsev

*Division of Theoretical Physics,
Yaroslavl State (P.G. Demidov) University,
Sovietskaya 14, 150000 Yaroslavl, Russian Federation
E-mail: mch@uniyar.ac.ru, rda@uniyar.ac.ru*

Abstract

The process of Compton scattering $\gamma e^\pm \rightarrow \gamma e^\pm$ in strongly magnetized hot electron-positron plasma is considered. The analytical expressions for the partial cross sections in rarefied plasma are obtained. The numerical estimations of the absorption rates for various scattering channels are presented with taking into account of the photon dispersion and wave function renormalization in strong magnetic field and plasma. The comparison of the scattering absorption rate with photon splitting probability shows the existence of the temperature range where these values are comparable with each other. The astrophysical applications of the obtaining results are discussed.

1 Introduction

Recent observations [1–4] give ground to believe that some astrophysical objects (SGR and AXP) are magnetars, a distinct class of isolated neutron stars with magnetic field strength of $B \sim 10^{14} - 10^{16}$ G [5–7], i.e. $B \gg B_e$, where $B_e = m^2/e \simeq 4.41 \times 10^{13}$ G¹ is the critical magnetic field. The spectra analysis of these objects is also providing evidence for the presence of electron-positron plasma in magnetar environment. It is well-known that strong magnetic field and/or plasma could influence essentially on different quantum processes [8–11]. Therefore the effect of magnetized plasma on microscopic physics should be incorporated in the magnetosphere models of strongly magnetized neutron stars.

The various studies indicate that electromagnetic processes such as Compton scattering and photon splitting $\gamma \rightarrow \gamma\gamma$ (merging $\gamma\gamma \rightarrow \gamma$) could play a crucial role in these models. For example, the process of photon splitting in strong magnetic field could suppress the production of electron-positron pair required for a detectable radio emission from magnetar [12, 13]. In turn, Compton scattering may play an important role in the models of atmosphere emission from strongly magnetized neutron stars [14, 15]. The both processes are very important in the models the radiation emission from SGR burst [6, 16, 17].

Previously, Compton scattering was studied under various conditions. The case of the magnetic field without plasma was investigated in [18–21]. It was found that Compton scattering becomes strongly anisotropic and essentially depends on photon polarization. In [15] it was

¹We use natural units $c = \hbar = k = 1$, m is the electron mass, $e > 0$ is the elementary charge.

shown that the dispersion properties of photon in strongly magnetized cold plasma could significantly influence on the process under consideration.

In the present work the process of photon scattering, $\gamma e^\pm \rightarrow \gamma e^\pm$, is investigated in the presence of strong magnetic field and charge-symmetric electron positron plasma, when the magnetic field strength B is the maximal physical parameter, namely $\sqrt{eB} \gg T, \omega, E$. Here T is the plasma temperature, ω and E is the initial photon and electron energies. In this case almost all electrons and positrons in plasma are on the ground Landau level. We also consider the none resonance case, when electron and photon momenta satisfy the condition $eB \gg (pk)$. Such conditions could be realized e.g. in the Thompson & Duncan model of SGR burst [6] in which magnetically trapped high temperature (~ 1 MeV) plasma fireball could be created.

The main purposes of our work are:

- to investigate the influence of the strongly magnetized hot plasma on the process of Compton scattering taking account of photon dispersion and wave function renormalization;
- to compare the Compton scattering with the process of photon splitting;
- to estimate the possible effect of the process under consideration on spectral formation of SGR.

The plan of the paper is following. In the section 2 we consider the propagation of the electromagnetic radiation in magnetized medium and give the expression of Compton scattering amplitudes for different polarization configurations of the initial and final photons. The analytic and numerical calculations of the photon absorption rate and cross section of the process are presented in Section 3. The discussion of the possible astrophysical application of Compton scattering are given in Section 4.

2 Photon propagation in strongly magnetized hot plasma

The propagation of the electromagnetic radiation in any active medium is convenient to describe in terms of normal modes (eigenmodes). In turn, the polarization and dispersion properties of normal modes are connected with eigenvectors and eigenvalues of polarization operator correspondingly. In the case of strongly magnetized charge-symmetric plasma the eigenvalues of the polarization operator can be obtained from the results of [22]:

$$\mathcal{P}^{(1)}(q) \simeq -\frac{\alpha}{3\pi} q_\perp^2 - q^2 \Lambda(B), \quad (1)$$

$$\mathcal{P}^{(2)}(q) \simeq -\frac{2eB\alpha}{\pi} \left[H\left(\frac{q_\parallel^2}{4m^2}\right) + \mathcal{J}(q_\parallel) \right] - q^2 \Lambda(B), \quad (2)$$

$$\mathcal{P}^{(3)}(q) \simeq -q^2 \Lambda(B), \quad (3)$$

where

$$\Lambda(B) = \frac{\alpha}{3\pi} [1.792 - \ln(B/B_e)].$$

$$\mathcal{J}(q_\parallel) = 4q_\parallel^2 m^2 \int \frac{dp_z}{E} \frac{f_E}{(q_\parallel^2)^2 - 4(pq_\parallel)^2}, \quad E = \sqrt{p_z^2 + m^2},$$

$f_E = [\exp(E/T) + 1]^{-1}$ is the electron distribution function,

$$H(z) = \frac{1}{\sqrt{z(1-z)}} \arctan \sqrt{\frac{z}{1-z}} - 1, \quad z \leq 1.$$




Figure 1: Photon dispersion laws in strong magnetic field $B/B_e = 200$ and neutral plasma vs. temperature: $T = 1$ MeV (is upper curve), $T = 0.5$ MeV (is middle curve), $T = 0.25$ MeV (is lower curve). Photon dispersion without plasma is depicted by dashed line. Dotted line corresponds to the vacuum dispersion law, $q^2 = 0$. The angle between the photon momentum and the magnetic field direction is $\pi/2$.

The four-vectors with indices \perp and \parallel belong to the Euclidean $\{1, 2\}$ -subspace and the Minkowski $\{0, 3\}$ -subspace correspondingly in the frame where the magnetic field is directed along third axis; $\varphi_{\alpha\beta} = F_{\alpha\beta}/B$ and $\tilde{\varphi}_{\alpha\beta} = \frac{1}{2}\varepsilon_{\alpha\beta\mu\nu}\varphi_{\mu\nu}$ are the dimensionless field tensor and dual field tensor correspondingly. The tensors $\Lambda_{\alpha\beta} = (\varphi\varphi)_{\alpha\beta}$, $\tilde{\Lambda}_{\alpha\beta} = (\tilde{\varphi}\tilde{\varphi})_{\alpha\beta}$, with equation $\tilde{\Lambda}_{\alpha\beta} - \Lambda_{\alpha\beta} = g_{\alpha\beta} = \text{diag}(1, -1, -1, -1)$ are introduced.

The dispersion properties of the normal modes are defined from the dispersion equations

$$q^2 - \mathcal{P}^{(\lambda)}(q) = 0 \quad (\lambda = 1, 2, 3). \quad (4)$$

Their analysis shows that the modes with $\lambda = 1, 2$ and with polarization vectors

$$\varepsilon_{\alpha}^{(1)}(q) = \frac{(q\varphi)_{\alpha}}{\sqrt{q_{\perp}^2}}, \quad \varepsilon_{\alpha}^{(2)}(q) = \frac{(q\tilde{\varphi})_{\alpha}}{\sqrt{q_{\parallel}^2}}. \quad (5)$$

are only physical in the case under consideration, just as in the pure magnetic field ². However, it should be emphasized that the coincidence of the polarization vectors in the plasma is approximate, to within $O(1/\beta)$.

Notice, that in plasma only the eigenvalue $\mathcal{P}^{(2)}(q)$ modifies in comparison with pure magnetic field case. It means that the dispersion law of the mode 1 is the same one as in the magnetized vacuum, where its deviation from the vacuum law, $q^2 = 0$, is negligibly small. From the other hand, the dispersion properties of the mode 2 essentially differs from the magnetized vacuum ones. In the Figure 1 the photon dispersion in both strong magnetic field and magnetized plasma at various temperatures are depicted. One can see that in the presence of the magnetized plasma there exist the kinematical region, where $q^2 > 0$ contrary to the case of pure magnetic field. This fact could lead to the alteration of the different photon processes kinematic. For example, the photon splitting channel $\gamma_2 \rightarrow \gamma_1\gamma_1$ forbidden in the magnetic field without plasma becomes allowed [24]. Another important distinction is an essentially different dependence of the dispersion law in variables q_{\parallel}^2 and q_{\perp}^2 on the angle between the photon momentum and the magnetic field direction (see Fig. 2).

²Symbols 1 and 2 correspond to the \parallel and \perp polarizations in pure magnetic field [23] and E - and O - modes in magnetized plasma [6].




Figure 2: Photon dispersion in a strong magnetic field ($B/B_e = 200$) and a charge-symmetric plasma ($T = 1$ MeV) at various angles between the photon momentum and the magnetic field direction: $\theta = \pi/2$ (upper solid curve), $\pi/6$ (middle solid curve), and $\pi/12$ (lower solid curve). The dashed curve represents the dispersion in the absence of a plasma.

It follows from Eq. (2) that the eigenvalue of the polarization operator $\mathcal{P}^{(2)}$ becomes large near the electron- positron pair production threshold, $q_{\parallel}^2 = 4m^2$. This suggests that the renormalization of the wave function for a photon of this polarization should be taken into account:

$$\varepsilon_{\alpha}^{(2)}(q) \rightarrow \varepsilon_{\alpha}^{(2)}(q)\sqrt{Z_2}, \quad Z_2^{-1} = 1 - \frac{\partial \mathcal{P}^{(2)}(q)}{\partial \omega^2}. \quad (6)$$

To calculate the amplitude of process in strong magnetic field one should use the Dirac equation solutions at the ground Landau level. For the electron propagator it is relevant to use its asymptotic form [10]. Then substituting the polarization vectors (5) one can obtain partial amplitudes for different polarization configuration of the initial and final photons. It is possible to present them in the covariant form:

$$\mathcal{M}_{1 \rightarrow 1} = -\frac{8\pi\alpha m}{eB} \frac{(q\varphi q')(q\tilde{\varphi}q')}{\sqrt{q_{\perp}^2 q_{\perp}'^2 (-Q_{\parallel}^2)}}, \quad (7)$$

$$\mathcal{M}_{1 \rightarrow 2} = -\frac{8\pi\alpha m}{eB} \frac{(q\Lambda q')(q'\tilde{\Lambda}Q)}{\sqrt{q_{\perp}^2 q_{\parallel}'^2 (-Q_{\parallel}^2)}}, \quad (8)$$

$$\mathcal{M}_{2 \rightarrow 1} = \frac{8\pi\alpha m}{eB} \frac{(q\Lambda q')(q\tilde{\Lambda}Q)}{\sqrt{q_{\parallel}^2 q_{\perp}'^2 (-Q_{\parallel}^2)}}, \quad (9)$$

$$\mathcal{M}_{2 \rightarrow 2} = 16i\pi\alpha m \frac{\sqrt{q_{\parallel}^2 q_{\parallel}'^2} \sqrt{(-Q_{\parallel}^2)} \varkappa}{(q\tilde{\Lambda}q')^2 - \varkappa^2 (q\tilde{\varphi}q')^2}, \quad (10)$$

where $\varkappa = \sqrt{1 - 4m^2/Q_{\parallel}^2}$ and $Q_{\parallel}^2 = (q - q'_{\parallel})^2 < 0$. We can see from last equations that all amplitudes except $\mathcal{M}_{2 \rightarrow 2}$ are suppressed by magnetic field strength. Therefore one could expect that mode 2 has the largest scattering absorption rate.

3 Photon absorption rate and cross section in the strongly magnetized medium

To analyse the efficiency of the process under consideration and to compare it with other competitive reactions we calculate the photon absorption rate which can be defined in the following way:

$$W_{\lambda e^{\pm} \rightarrow \lambda' e^{\pm}} = \frac{eB}{16(2\pi)^4 \omega_{\lambda}} \int |\mathcal{M}_{\lambda \rightarrow \lambda'}|^2 Z_{\lambda} Z_{\lambda'} \times \quad (11)$$

$$\times f_E (1 - f_{E'}) (1 + f_{\omega'}) \delta(\omega_{\lambda}(\mathbf{k}) + E - \omega_{\lambda'}(\mathbf{k}') - E') \frac{dp_z d^3 k'}{EE' \omega_{\lambda'}},$$

where $f_{\omega'} = [\exp(\omega'/T) - 1]^{-1}$ is the photon distribution function. In the case of the rarefied plasma ($T \ll m$) these absorption rates can be expressed in term of partial cross sections³ $W_{\lambda \rightarrow \lambda'} = W_{\lambda e^- \rightarrow \lambda' e^-} + W_{\lambda e^+ \rightarrow \lambda' e^+} = n_e \sigma_{\lambda \rightarrow \lambda'}$:

$$\sigma_{1 \rightarrow 1} = \frac{3}{8} \sigma_T \left(\frac{B_e}{B} \right)^2 \frac{\omega^2}{m^2} \left[\frac{\omega + 2m}{2(\omega + m)} + \frac{m}{\omega} \ln \left(1 + \frac{\omega}{m} \right) \right], \quad (12)$$

$$\sigma_{2 \rightarrow 1} = \frac{3}{8} \sigma_T \left(\frac{B_e}{B} \right)^2 \frac{q_{\perp}^2}{m^2} Z_2 \left[\frac{\omega + 2m}{2(\omega + m)} - \frac{m}{\omega} \ln \left(1 + \frac{\omega}{m} \right) \right], \quad (13)$$

$$\sigma_{1 \rightarrow 2} = \frac{3}{4} \sigma_T \left(\frac{B_e}{B} \right)^2 \frac{(\omega + 2m)^2}{\omega(\omega + m)} \int_0^{\omega^2/4m^2} dz \left(1 + \frac{3}{2} \xi \frac{H(z)}{z} \right) \sqrt{\frac{\omega^2 - 4m^2 z}{(\omega + 2m)^2 - 4m^2 z}}, \quad (14)$$

$$\sigma_{2 \rightarrow 2} = 6 \sigma_T \frac{m^4(\omega + m)}{\omega^3(\omega + 2m)^2} Z_2 \left[\frac{\omega(\omega + 2m)}{(\omega + m)(2m - \omega)} - \ln \left(1 + \frac{\omega}{m} \right) + \frac{2\omega(\omega - m)(2m + \omega)}{(\omega + m)(2m - \omega)\sqrt{4m^2 - \omega^2}} \arctan \frac{\omega}{\sqrt{4m^2 - \omega^2}} \right], \quad (15)$$

where σ_T is the Thompson cross section, $\xi = \frac{\alpha B}{3\pi B_e}$ is the parameter characterizing magnetic field influence, $q_{\perp}^2 = \omega^2 - \mathcal{P}^{(2)}(\omega^2)$ and the number of electron (positron) density in a strongly magnetized, charge-symmetric rarefied plasma can be estimated as

$$n_e \simeq eB \sqrt{\frac{mT}{2\pi^3}} e^{-m/T}. \quad (16)$$

In the low energy limit ($\omega \ll m$) the formulas (12-15) can be presented in the following form:

$$\sigma_{1 \rightarrow 1} = \frac{3}{4} \sigma_T \left(\frac{B_e}{B} \right)^2 \frac{\omega^2}{m^2}, \quad \sigma_{1 \rightarrow 2} = \frac{1}{4} \sigma_T \left(\frac{B_e}{B} \right)^2 \frac{\omega^2}{m^2} (1 + \xi), \quad (17)$$

$$\sigma_{2 \rightarrow 1} = \frac{1}{16} \sigma_T \left(\frac{B_e}{B} \right)^2 \frac{\omega^2(\omega^2 - \omega_{pl}^2)}{m^4} \theta(\omega - \omega_{pl}), \quad \sigma_{2 \rightarrow 2} = \frac{\sigma_T}{1 + \xi}, \quad (18)$$

where ω_{pl} is plasma frequency defined by equation $\omega_{pl}^2 - \mathcal{P}^{(2)}(\omega_{pl}, \vec{k} \rightarrow 0) = 0$. One can see that the presence of magnetized plasma slightly influences on the process cross-sections in this limit. Moreover, the corrections connected with photon dispersion and wave function renormalization

³Hereafter the initial photon propagation across magnetic field is considered




Figure 3: The dependence of the photon absorption rate for channels $\gamma_2 e^\pm \rightarrow \gamma_2 e^\pm$ (solid line) and $\gamma_2 e^\pm \rightarrow \gamma_1 e^\pm$ (dashed line) on energy of initial photon in strong magnetic field $B/B_e = 200$ at $T = 1\text{MeV} - a$, $T = 250\text{keV} - b$ and $T = 50\text{keV} - c$. The dotted line corresponds to the probability of photon splitting, $\gamma_2 \rightarrow \gamma_1 \gamma_1$, at $T = 1\text{MeV}$ [24]. Here $W_0 = (\alpha/\pi)^3 m \simeq 3.25 \cdot 10^2 \text{cm}^{-1}$.

are significant only for $\xi \sim 1$, i.e. when magnetic field is rather strong $B \sim 10^3 B_e$. In the case $\xi \ll 1$ which is relevant for the models of magnetar magnetosphere emission the formulas (17, 18) coincide with the well-known result [18]. It is interesting also to consider the case of $\omega \sim 2m$. In this region the magnetized vacuum influence is large and the cross-sections (12-15) have the form:

$$\sigma_{1 \rightarrow 1} = \sigma_T \left(\frac{B_e}{B}\right)^2 \left[1 + \frac{3}{4} \ln 3\right], \quad \sigma_{1 \rightarrow 2} = 4\sigma_T \left(\frac{B_e}{B}\right)^2 \left[1 - \frac{3}{4} \ln 3\right] (1 + 2\xi), \quad (19)$$

$$\sigma_{2 \rightarrow 1} = 2\sigma_T \left(\frac{B_e}{B}\right)^2 \left[1 - \frac{3}{4} \ln 3\right] \left(1 - \frac{\omega^2}{4m^2}\right), \quad \sigma_{2 \rightarrow 2} = \frac{\sigma_T}{2\xi}. \quad (20)$$

To investigate the Compton scattering under hot plasma conditions ($T \sim m$) we have made the numerical calculations of photon absorption rates for the various channels. The results are represented in Figures 3 and 4.

4 Discussion

In the models of soft gamma repeaters spectrum formation the dependence of photon absorption rates on energy and temperature plays an important role. It could influence on the shape of emergent spectrum and defines the temperature profile in the emission region during bursts in SGRs [6, 25]. In the both figures 3 and 4 one can see that photon absorption rates corresponding to Compton scattering are the fast increasing functions of temperature. At the same time, the channels with initial photon of mode 1 and mode 2 have different character of the absorption coefficient energy dependence. As shown in the Fig.3 the absorption rates for the reactions $\gamma_2 e^\pm \rightarrow \gamma_2 e^\pm$ and $\gamma_2 e^\pm \rightarrow \gamma_1 e^\pm$ have thresholds. They are caused by mode 2 dispersion relation with plasma frequency and indicated the fact that the electromagnetic wave corresponding to mode 2 can't propagate with energy below ω_{pl} . In the region above plasma frequency $W_{\gamma_2 e^\pm \rightarrow \gamma_2 e^\pm}$ is slightly depends on energy while $W_{\gamma_2 e^\pm \rightarrow \gamma_1 e^\pm}$ tends to zero in the vicinity of ω_{pl} and pair creation threshold, $\omega = 2m$. We would like to note that in the vicinity of pair creation threshold taking account of wave function renormalization and photon




Figure 4: The dependence of the photon absorption rate for channels $\gamma_1 e^\pm \rightarrow \gamma_1 e^\pm$ (solid line) and $\gamma_1 e^\pm \rightarrow \gamma_2 e^\pm$ (dashed line) on energy of initial photon in strong magnetic field $B/B_e = 200$ at $T = 1\text{MeV}$ - *a*, $T = 250\text{keV}$ - *b* and $T = 50\text{keV}$ - *c*. The dotted and chain lines correspond to the probability of photon splitting, $\gamma_1 \rightarrow \gamma_1\gamma_2$ and $\gamma_1 \rightarrow \gamma_2\gamma_2$, respectively, at $T = 50\text{keV}$ [24]. Here $W_0 = (\alpha/\pi)^3 m \simeq 3.25 \cdot 10^2 \text{cm}^{-1}$.

dispersion becomes very important and defines the processes rates dependence on energy, temperature and magnetic field. In the region $\omega \geq 2m$ the photon of mode 2 is unstable due to the process $\gamma_2 \rightarrow e^+e^-$ with rate much larger than Compton scattering. The energy dependence of $W_{\gamma_1 e^\pm \rightarrow \gamma_1 e^\pm}$ and $W_{\gamma_1 e^\pm \rightarrow \gamma_2 e^\pm}$ is depicted in Fig.4. One can see the fast increasing of absorption coefficients at low energies and rather slow dependence at $\omega \gtrsim 2m$.

The previous investigation of radiation transfer problem in strongly magnetized plasma have shown that along with Compton scattering process the photon splitting $\gamma \rightarrow \gamma\gamma$ could play a significant role as mechanism of photon production [6]. To compare these processes we depict the probabilities of the photon splitting channels on the same plots (see dotted and chain lines in Fig.3 and Fig.4). One can see that at temperature $T \sim m$ and in kinematical region $\omega \leq 2m$ the main photon splitting process is $\gamma_2 \rightarrow \gamma_1\gamma_1$ forbidden in pure magnetic field [24]. It is seen also that the rate of the process is much less than Compton scattering ones. Nevertheless, it could be an effective photon production mechanism at temperatures under consideration. The probabilities of the channels $\gamma_1 \rightarrow \gamma_1\gamma_2$ and $\gamma_1 \rightarrow \gamma_2\gamma_2$ increase with temperature falling and become comparable and even larger than Compton scattering rates $W_{\gamma_1 e^\pm \rightarrow \gamma_1 e^\pm}$ and $W_{\gamma_1 e^\pm \rightarrow \gamma_2 e^\pm}$. As shown in Fig. 4 the process of photon splitting strongly dominates over Compton scattering at $T = 50\text{keV}$. It was claimed previously that the effect of the cold strongly magnetized plasma on photon splitting is not pronounced and the vacuum approximation can be used in the most calculations [15, 26]. We can see now that in the presence of hot plasma the process of photon splitting could not be only intensive source of photon production but also an effective absorption mechanism.

Let us illustrate this fact in the framework of the magnetar model of SGR burst. It is known that the radiation transfer in the magnetically trapped plasma may be described as a diffusion of the 1-mode photons whereas 2-mode photons are locked [6, 25]. The last circumstance is connected especially with weak dependence of the 2-mode absorption coefficient on photon energy (see Fig. 3). In the assumption that the properties of the radiation field do not change much over absorption length W^{-1} the radiation transfer may be conveniently described in terms




Figure 5: The ratio of the Rosseland mean free paths without and with taking into account of photon splitting process as a function of the inverse temperature at $B = 200B_e$

of the Rosseland mean free path of 1-mode photons [6, 25]

$$\lambda_R = \frac{\int W^{-1} \omega^4 \exp(\omega/T) [\exp(\omega/T) - 1]^{-2} d\omega}{\int \omega^4 \exp(\omega/T) [\exp(\omega/T) - 1]^{-2} d\omega}, \quad (21)$$

where W is the total absorption coefficient. Then the temperature distribution in the magnetically trapped plasma region could be defined from the diffusion equation:

$$F(z) = \frac{16\lambda_R \sigma}{3} T^3 \frac{\partial T}{\partial z}, \quad (22)$$

where σ is the Stefan-Boltzmann constant, $F(z)$ is the radiation flux. If only Compton scattering process is considered then

$$W = W_{\gamma_1 e^\pm \rightarrow \gamma_1 e^\pm} + W_{\gamma_1 e^\pm \rightarrow \gamma_2 e^\pm}. \quad (23)$$

To take into account the effect of photon splitting one should add term $W_{\gamma_1 \rightarrow \gamma_1 \gamma_2} + W_{\gamma_1 \rightarrow \gamma_2 \gamma_2}$ to the equation (23). In the figure 5 the ratio of the Rosseland mean free path for only Compton scattering to Rosseland mean free path with photon splitting is depicted. One can see that at low temperatures the additional absorption process of photon splitting leads to the significant decreasing of the Rosseland mean free path. The solution of the diffusion equation (22) is out of the scope of this article. However, we would like to note that the more detail analysis of the radiation transfer needs the consistent solution of Boltzmann equation for the photon occupation number and radiative transfer equation in the wide range of temperatures ($10keV \lesssim T \lesssim 1MeV$).

In conclusion, we have investigated the influence of the strongly magnetized hot plasma on the Compton scattering process with taking into account of photon dispersion and large radiative corrections. It was found that absorption rate of the dominant scattering channel of the 2-mode photon weakly depends on energy in wide temperature range whereas the 1-mode absorption coefficient is fast increasing function of photon energy. The comparison of the photon splitting effect and Compton scattering shows that the influences of these reaction on the 1-mode radiation transfer are competitive in rarefied plasma ($T \ll m$).

Acknowledgements

We express our deep gratitude to the organizers of the Seminar ‘‘Quarks-2006’’ for warm hospitality. This work supported in part by the Council on Grants by the President of Russian Federation for the Support of Young Russian Scientists and Leading Scientific Schools of Russian Federation under the Grant No. NSh-6376.2006.2 and No. MK-1097.2005.2 (MC), and by the Russian Foundation for Basic Research under the Grant No. 04-02-16253. Work of MC is supported by the DAAD (Deutscher Akademischer Austauschdienst).

References

- [1] C. Kouveliotou *et al.*, Nature **393** (1998) 235.
- [2] C. Kouveliotou *et al.*, Astrophys. J. **510** (1999) L115
- [3] F. P. Gavriil, V. M. Kaspi, P. M. Woods, Nature **419** (2002) 142.
- [4] A. I. Ibrahim, S. Safi-Harb, J. H. Swank, W. Parke and S. Zane, Astrophys. J. **574** (2002) L51
- [5] R.C. Duncan, C. Thompson, Astrophys. J. **392** No. 1. (1992) L9
- [6] C. Thompson, R.C. Duncan, Mon. Not. R. Astron. Soc. **275** (1995) 255
- [7] R.C. Duncan, C. Thompson, Astrophys. J. **473** (1996) 322
- [8] R. C. Duncan, Fifth Huntsville Gamma-Ray Burst Symposium, Eds: R. Marc Kippen, Robert S. Mallozzi, Gerald J. Fishman. AIP Vol. 526 (American Institute of Physics, Melville, New York) p.830 [astro-ph/0002442].
- [9] D. Lai, Rev. Mod. Phys. **73** (2001) 629
- [10] A. V. Kuznetsov, N. V. Mikheev, *Electroweak Processes in External Electromagnetic Fields*, Springer-Verlag, New York (2003), p.1.
- [11] A. K. Harding, D. Lai, Rept. Prog. Phys. **69** (2006) 2631
- [12] A.K. Harding, M.G. Baring, Astrophys. J. **507** (1998) L55
- [13] M. G. Baring and A. K. Harding, Astrophys. J. **547** (2001) 929
- [14] M.C. Miller, Astrophys. J. Lett. **448** (1995) L29
- [15] T. Bulik, M.C. Miller, Mon. Not. R. Astron. Soc. **288** (1997) 596
- [16] A. K. Harding, M. G. Baring, P. L. Gonthier, Astro. Astrophys. Suppl. Ser. **120** (1996) 111
- [17] A. K. Harding, M. G. Baring, P. L. Gonthier, Astrophys. J. **476** (1997) 246
- [18] H. Herold, Phys. Rev. **D19** (1979) 2868
- [19] D.B. Melrose, A.J. Parle, Aust. J. Phys. **36** (1983) 799
- [20] J.K. Daugherty, A.K. Harding, Astrophys. J. **309** (1986) 362
- [21] P.L. Gonthier, A.K. Harding, M.G. Baring et al., Astrophys. J. **540** (2000) 907
- [22] A.E. Shabad, Tr. Fiz. Inst. Akad. Nauk SSSR **192** (1988) 5
- [23] S. L. Adler, Ann. Phys. (N.Y.) **67** (1971) 599
- [24] D. A. Romyantsev, M. V. Chistyakov, JETP **101** (2005) 635
- [25] Y. E. Lyubarsky, Mon. Not. R. Astron. Soc. **332** (2002) 199
- [26] P Elmfors., B Skagerstam., Phys. Lett. **B427** (1998) 197